



NATIONAL SENIOR CERTIFICATE

GRADE 12

SEPTEMBER 2024

MATHEMATICS P1

MARKS: 150

TIME: 3 hours



This question paper consists of 13 pages, including a 1-page information sheet.

INSTRUCTIONS AND INFORMATION

Read the following instructions carefully before answering the questions.

1. This question paper consists of 12 questions.
2. Answer ALL the questions.
3. Clearly show ALL calculations, diagrams, graphs, et cetera which you have used in determining your answers.
4. You may use an approved scientific calculator (non-programmable and non-graphical), unless stated otherwise.
5. Answers only will NOT necessarily be awarded full marks.
6. If necessary, answers should be rounded off to TWO decimal places, unless stated otherwise.
7. Diagrams are NOT necessarily drawn to scale.
8. Number the answers correctly according to the numbering system used in this question paper.
9. An information sheet with formulae is included at the end of the question paper.
10. Write neatly and legibly.



QUESTION 1

1.1 Solve for x :

1.1.1 $(2x-4)(x-1)=0$ (2)

1.1.2 $2x^2-3(x+2)=4$ (correct to TWO decimal places) (4)

1.1.3 $x^2+4x-21\leq 0$ (3)

1.1.4 $-\sqrt{x-1}=3-2x$ (4)

1.2 Solve simultaneously for x and y :

$2x=1-y$ and $xy-x^2+y^2=5$ (6)

1.3 Given that:

- $f(x)=x^2+3x$
- $2x-[t(x)]^{\frac{1}{2}}=0$

For which values of k will the equation $f(-x)+\frac{t(2k)}{4}=0$ have equal roots? (5)

[24]



QUESTION 2

2.1 Given the quadratic number pattern: $-5; -4; -1; 4; \dots$

2.1.1 Determine the n^{th} term of the quadratic number pattern in the form $T_n = an^2 + bn + c$. (4)

2.1.2 Calculate the 35th term of the quadratic number pattern. (1)

2.1.3 Which TWO consecutive terms of the first differences sequence will have a product of 1 155? (4)

2.2 Given the arithmetic sequence: $60; 65; 70; \dots$

Calculate the value of p for which $T_p = 430$. (3)

2.3 The sum of the first three terms of an increasing arithmetic series is 30 and the product of the same three terms is 510. Determine the values of a and d , the first term and the common difference of the series respectively. (5)

[17]

QUESTION 3

3.1 An infinite geometric series has a first term of 2 and constant ratio of $\frac{1}{3}$.

3.1.1 Calculate the next two terms. (1)

3.1.2 Calculate the value of S_{∞} . (2)

3.2 Determine the value of m if:

$$\sum_{k=3}^m 8(2)^{k-1} = 131\,040$$


(5)
[8]



QUESTION 4


Consider the function: $f(x) = \frac{-1}{x+5} - 2$

- 4.1 Write down the equations of asymptotes of f . (2)
- 4.2 Determine the coordinates of the x -intercept of f . (2)
- 4.3 Determine the coordinates of the y -intercept of f . (2)
- 4.4 Sketch the graph of f , show clearly all asymptotes and intercepts with the axes. (3)
- 4.5 Determine the equation of the axis of symmetry that has a gradient of -1 . (2)

[11]

QUESTION 5

The graphs of $f(x) = x^2 - 5x + 6$ and $g(x) = x + 1$ are drawn below. B and C are points of intersection of f and g . The graph of f has x -intercepts at $(2;0)$ and $(3;0)$ and a y -intercept at A.



- 5.1 Determine the equation of the axis of symmetry of f . (2)
- 5.2 Calculate the coordinates of B and C. (4)
- 5.3 PQ is the vertical distance between the graphs g and f between B and C. Determine the maximum length of PQ. (4)
- 5.4 Determine the range of $t(x)$ if $f(x) - 2 = t(x)$. (2)
- 5.5 For which values of x is $f(x) \cdot g'(x) < 0$? (2)


[14]



QUESTION 6

The diagram below shows the graphs of $f(x) = -\log_c x$ and $g(x) = dx^2$; $x \geq 0$.

The point $P\left(\frac{1}{2}; \frac{1}{2}\right)$ is the point of intersection of the graphs f and g .



6.1 Calculate the values of c and d . (3)

6.2 Determine:

6.2.1 The equation of $g^{-1}(x)$ in the form $y = \dots$ (2)

6.2.2 The equation of $h^{-1}(x)$ in the form $y = \dots$, if h is a reflection of f in the x -axis (2)

6.2.3 The x -values for which $h^{-1}(x) > 0$ (1)
[8]



QUESTION 7

- 7.1 A car that is worth R180 000, depreciates at 13% p.a. compounded annually on the reducing balance method. Calculate the value of the car in 6 years. (3)

- 7.2 Lumi opened a 15-year savings plan account that pays interest at 8% per annum compounded monthly. She saves an amount of R900 every month for the first 10 years. Her first payment was at the end of the first month. For the last 5 years of her savings plan she managed to increase her monthly payments to R1 300.

Calculate the value of her savings at the end of the savings period. (5)

- 7.3 Mr Leanya bought a house for R850 000. He obtained a loan from the bank at an interest rate of 13% per annum compounded monthly to pay for the house. He agreed to pay monthly instalments of R9 958,39 for 20 years.

- 7.3.1 Calculate the balance of his loan immediately after his 75th instalment. (3)

- 7.3.2 Mr Leanya experienced financial difficulties after his 75th instalment and did not pay the 76th to the 79th instalments. At the end of the 80th month he increased his monthly instalment so as to pay off the loan in the same time interval as planned initially.

Calculate the value of his new adjusted monthly instalment. (5)

[16]

QUESTION 8

- 8.1 Determine $f'(x)$ from first principles if $f(x) = x^2 - 3$. (4)

- 8.2 Determine:

8.2.1 $\frac{dy}{dx}$ if $y = -3x^2 + 7x$ (2)

8.2.2 $D_x \left[\frac{x^3 - 5x^2}{x^3} - \sqrt{x} \right]$ (4)


- 8.3 Suppose that $g(x)$ represents the rate of change of $h(x) = -x^3 - 3x^2 + 1$. Calculate the largest value of $g(x)$. (3)

[13]



QUESTION 9

- 9.1 The sketch below shows the graph of $f(x) = -3x^3 + mx^2 + nx$. The graph of f passes through the origin and has a local minimum and a local maximum at $A(1; -12)$ and $B(3; 0)$ respectively.



- 9.1.1 Show that $m = 18$ and $n = -27$ (5)
- 9.1.2 Explain the difference between $f(a)$ and $f'(a)$. (2)
- 9.1.3 $g(x)$ is the tangent to the curve of $f(x)$ at the point of inflection. Determine the equation of $h(x)$, the straight line that is perpendicular to $g(x)$ and passes through the origin. (5)
- 9.1.4 For which values of x will $f''(x) > 0$? (2)
- 9.2 The function t is defined by $t(x) = 2x^3 + bx + c$ and has the following properties.
- $t(-3) = t(3) = t(0) = 0$
 - $t'(-1, 5) = t'(1, 5) = 0$

Use this information to draw a neatly labelled sketch graph of t , without solving for b and c .


(3)
[17]

QUESTION 10

The number of scripts marked by a certain marker was tracked at a marking centre t days after marking started, and is represented by the function, $S(t) = -3t^2 + 30t$, $1 \leq t \leq 10$, $t \in \mathbb{Z}$, where $S(t)$ is measured in scripts per day.


- 10.1 Determine the number of scripts that were marked by the marker on the third day. (2)
- 10.2 On which day will the marker reach the maximum number of scripts marked per day? (3)
- 10.3 The total number of scripts that a marker had to mark for the 10 days was 500. Did this marker reach the quota? Support your answer with calculations. (2)

[7]



QUESTION 11

- 11.1 Two events A and B are shown on the Venn diagram below.
It is given that $P[\text{not } (A \text{ or } B)] = 0,41$.



Determine:

- 11.1.1 The value of x and hence $P(A)$ (2)
- 11.1.2 $P(A \text{ or not } B)$ (2)
- 11.2 The results for the soccer club, City Brothers FC's 30 games during the 2022–2023 season are shown below.

	HOME GAME	AWAY GAME	TOTAL
WINS	3	4	7
LOSSES	7	7	14
DRAWS	5	a	9
TOTAL	15	15	30

- 11.2.1 Write down the value of a . (1)
- 11.2.2 What is the probability that in a randomly selected match City Brothers FC was the losing team? (1)
- 11.2.3 Are the events 'winning a game' and 'playing at the home ground' independent? Justify your answer with calculations. (3)

[9]



QUESTION 12

The province of KwaZulu-Natal has introduced a new number plate system starting from December 2023. The new number plate code consists of two letters, two digits and then two letters. The system is using the digits, 0–9 and the letters of the alphabet excluding the vowels. Below is an example of this new number plate. Note that all number plates come with the ZN suffix which is independent from the code.



[Source: KZN Provincial Gazette 2614-new number plates for KZN]

- 12.1 How many number plate codes are possible with the new system, if digits and letters may not be repeated? (2)
- 12.2 Calculate the probability that a number plate code will start with a letter of the alphabet that is before letter G, with the first digit being a composite number and that the last digit is a factor of 4. Digits and letters may not be repeated. (4)

[6]

TOTAL: 150



INFORMATION SHEET: MATHEMATICS

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$A = P(1 + ni)$$

$$A = P(1 - ni)$$

$$A = P(1 - i)^n$$

$$A = P(1 + i)^n$$

$$T_n = a + (n-1)d \quad S_n = \frac{n}{2}(2a + (n-1)d)$$

$$T_n = ar^{n-1} \quad S_n = \frac{a(r^n - 1)}{r - 1} ; \quad r \neq 1 \quad S_\infty = \frac{a}{1 - r} ; \quad -1 < r < 1$$

$$F = \frac{x[(1+i)^n - 1]}{i} \quad P = \frac{x[1 - (1+i)^{-n}]}{i}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$$

$$y = mx + c$$

$$y - y_1 = m(x - x_1)$$

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

$$m = \tan \theta$$

$$(x - a)^2 + (y - b)^2 = r^2$$

$$\text{In } \triangle ABC: \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cdot \cos A \quad \text{area } \triangle ABC = \frac{1}{2}ab \cdot \sin C$$

$$\sin(\alpha + \beta) = \sin \alpha \cdot \cos \beta + \cos \alpha \cdot \sin \beta$$

$$\sin(\alpha - \beta) = \sin \alpha \cdot \cos \beta - \cos \alpha \cdot \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cdot \cos \beta - \sin \alpha \cdot \sin \beta$$

$$\cos(\alpha - \beta) = \cos \alpha \cdot \cos \beta + \sin \alpha \cdot \sin \beta$$

$$\cos 2\alpha = \begin{cases} \cos^2 \alpha - \sin^2 \alpha \\ 1 - 2\sin^2 \alpha \\ 2\cos^2 \alpha - 1 \end{cases}$$

$$\sin 2\alpha = 2 \sin \alpha \cdot \cos \alpha$$

$$\bar{x} = \frac{\sum x}{n}$$

$$\sigma^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}$$

$$P(A) = \frac{n(A)}{n(S)}$$


$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$

$$\hat{y} = a + bx$$

$$b = \frac{\sum (x - \bar{x})(y - \bar{y})}{\sum (x - \bar{x})^2}$$


Dear Mr West

RE: CORRECTION TO MARKING GUIDELINE (MATHEMATICS P1)

7.2	$F_v = \frac{x[(1+i)^n - 1]}{i}$  $= \frac{900 \left[\left(1 + \frac{0,08}{12}\right)^{120} - 1 \right]}{\frac{0,08}{12}} \left(1 + \frac{0,08}{12}\right)^{60} + \frac{1300 \left[\left(1 + \frac{0,08}{12}\right)^{60} - 1 \right]}{\frac{0,08}{12}}$ $= 245\,305,23 + 95\,519,91$ $= R340\,825,14$ was omitted in the original MG	✓ $n = 120$ and/en $i = \frac{8\%}{12}$ or / of $\frac{8}{1200}$ ✓ $n = 60$ in F_v ✓ substitution into F_v / vervang in F_v ✓ $\times \left(1 + \frac{0,08}{12}\right)^{60}$ ✓ answer / antwoord
-----	--	--

(5)

OR/OF (alternative method)

7.2	$F_v = \frac{x[(1+i)^n - 1]}{i}$ $= \frac{900 \left[\left(1 + \frac{0,08}{12}\right)^{180} - 1 \right]}{\frac{0,08}{12}} + \frac{400 \left[\left(1 + \frac{0,08}{12}\right)^{60} - 1 \right]}{\frac{0,08}{12}}$ $= 311\,434,40 + 29\,390,74$ $= R340\,825,14$	✓ $n = 180$ and/en $i = \frac{8\%}{12}$ or / of $\frac{8}{1200}$ ✓ substitution into F_v / vervang in F_v working with 900 ✓ $n = 60$ in F_v ✓ substitution into F_v / vervang in F_v working with 400 ✓ answer / antwoord 
-----	---	--

(5)

Regards

C.F. HERADIEN
(ECIM)



**NATIONAL
SENIOR CERTIFICATE/
NASIONALE
SENIORSERTIFIKAAT**

GRADE/GRAAD 12




SEPTEMBER 2024


**MATHEMATICS P1/WISKUNDE V1
MARKING GUIDELINE/NASIENRIGLYN**

MARKS/PUNTE: 150



This marking guideline consists of 22 pages./
Hierdie nasienriglyn bestaan uit 22 bladsye.





1.2	$2x = 1 - y$.....(1) $xy - x^2 + y^2 = 5$.....(2) $y = 1 - 2x$.....(3) Subst/Vervang (3) into/in (2) $x(1 - 2x) - x^2 + (1 - 2x)^2 = 5$ $x - 2x^2 - x^2 + 1 - 4x + 4x^2 - 5 = 0$ $x^2 - 3x - 4 = 0$ $(x - 4)(x + 1) = 0$ $x = 4$ or/of $x = -1$ For/Vir $x = 4$: $y = 1 - 2(4)$ $y = -7$ For/Vir $x = -1$: $y = 1 - 2(-1)$ $y = 3$ OR / OF $y = 1 - 2x$.....(1) $xy - x^2 + y^2 = 5$.....(2) $x = \frac{1}{2} - \frac{y}{2}$.....(3) Subst/Vervang (3) into/in (2) $y\left(\frac{1}{2} - \frac{y}{2}\right) - \left(\frac{1}{2} - \frac{y}{2}\right)^2 + y^2 = 5$ $\frac{y}{2} - \frac{y^2}{2} - \left(\frac{1}{4} - \frac{y}{2} + \frac{y^2}{4}\right) + y^2 = 5$ $\frac{y}{2} - \frac{y^2}{2} - \frac{1}{4} + \frac{y}{2} - \frac{y^2}{4} + y^2 = 5$ $\frac{y^2}{4} + y - \frac{21}{4} = 0$ $y^2 + 4x - 21 = 0$ $(y + 7)(y - 3) = 0$ $y = -7$ or/of $y = 3$ For/Vir $y = -7$: $x = 4$ For/Vir $y = 3$: $x = -1$	$\checkmark y = 1 - 2x$ $\checkmark$ substitution / vervanging $\checkmark$ standard form / standaardvorm $\checkmark$ factors / faktore $\checkmark$ x-values / x-waardes $\checkmark$ y-values / y-waardes OR/OF $\checkmark x = \frac{1}{2} - \frac{y}{2}$ $\checkmark$ substitution / vervanging $\checkmark$ standard form / standaardvorm $\checkmark$ factors / faktore $\checkmark$ y-values / y-waardes $\checkmark$ x-values / x-waardes
------------	---	--

(6)



QUESTION 2/VRAAG 2


2.1		
2.1.1	$2a = 2$ $\therefore a = 1$ $3a + b = 1$ $b = 1 - 3(1)$ $\therefore b = -2$ $\therefore T_n = n^2 - 2n - 4$	$a + b + c = -5$ $c = -5 - 1 + 2$ $\therefore c = -4$ $\checkmark a = 1$ $\checkmark b = -2$ $\checkmark c = -4$ $\checkmark T_n = n^2 - 2n - 4$ (4)
2.1.2	$T_n = n^2 - 2n - 4$ $T_{35} = (35)^2 - 2(35) - 4$ $= 1\,151$	$\checkmark$ answer / antwoord (1)
2.1.3	$T_n = 1 + (n-1)(2)$ $T_n = 2n - 1$ $T_{n+1} = 2(n+1) - 1$ $T_{n+1} = 2n + 1$ $T_n \times T_{n+1} = 1155$ $(2n-1)(2n+1) = 1155$ $4n^2 - 1 = 1155$ $4n^2 = 1156$ $n^2 = 289$ $n = \pm 17$ $\therefore n = 17, n \in \mathbb{N}$ $\therefore T_{17}$ and/en T_{18} will give a product of 1155 <i>sal 'n produk van 1 155 gee</i> OR/OF $T_n = 2n - 1$ $T_{n-1} = 2n - 3$ $(2n-1)(2n-3) = 1155$ $4n^2 - 8n - 1152 = 0$ $n^2 - 2n - 288 = 0$ $(n-18)(n+16) = 0$ $n = 18$ or/of $n \neq -16, n \in \mathbb{N}$ $n = 18$ and/en $n-1 = 17$ $\therefore T_{17}$ and/en T_{18} will give a product of 1155 <i>sal 'n produk van 1 155 gee</i>	$\checkmark T_n = 2n - 1$ $\checkmark (2n-1)(2n+1) = 1155$ $\checkmark$ standard form / <i>standaardvorm</i> $\checkmark n = 17$ and/en $n+1 = 18$ OR/OF $\checkmark T_n = 2n - 1$ $\checkmark (2n-1)(2n-3) = 1155$ $\checkmark$ standard form / <i>standaardvorm</i> $\checkmark n = 18$ and/en $n-1 = 17$ (4)




QUESTION 3/VRAAG 3

3.1.1	$\frac{2}{3} ; \frac{2}{9}$	✓ answer / <i>antwoord</i> (1)
3.1.2	$S_{\infty} = \frac{a}{1-r}$ $= \frac{2}{1-\frac{1}{3}}$ $= 3$	✓ substitution / <i>vervanging</i> ✓ answer / <i>antwoord</i> (2)
3.2	$\sum_{k=3}^m 8(2)^{k+1} = 131\,040$ $32 + 64 + 128 + \dots$ $r = 2$ $s_n = \frac{a(1-r^n)}{1-r}$ $131\,040 = \frac{32(2^n - 1)}{2 - 1}$ $\therefore 2^n - 1 = 4\,095$ $2^n = 4\,096$ $2^n = 2^{12} \quad \text{OR/OF } n = \log_2(4\,096)$ $\Rightarrow n = 12$ $n = m - 3 + 1$ $12 = m - 2$ $m = 14$	✓ value of a / <i>waarde van a</i> ✓ substitution / <i>vervanging</i> ✓ simplification / <i>vereenvoudiging</i> ✓ value of n / <i>waarde van n</i> ✓ answer / <i>antwoord</i> (5)
		[8]

QUESTION 4/VRAAG 4

4.1	$x = -5$ $y = -2$	✓ equation of V.A / <i>vergelyking van V.A</i> ✓ equation of H.A / <i>vergelyking van H.A</i> (2)
4.2	$0 = \frac{-1}{x+5} - 2$ $2 = \frac{-1}{x+5}$ $2(x+5) = -1$ $2x = -11$ $x = \frac{-11}{2} = -5,5$ $\therefore \left(-\frac{11}{2}; 0\right)$	✓ $y = 0$ ✓ answer / <i>antwoord</i> (2)
4.3	$y = \frac{-1}{x+5} - 2$ $y = \frac{-1}{0+5} - 2$ $y = -\frac{11}{5} = -2,2$ $\therefore \left(0; -\frac{11}{5}\right)$	✓ $x = 0$ ✓ answer / <i>antwoord</i> (2)
4.4		✓ both asymptotes / <i>beide asimptote</i> ✓ x-intercept / <i>x-afsnit</i> and/of y-intercept / <i>y-afsnit</i> ✓ shape / <i>vorm</i> (3)
4.5	$y = -(x+5) - 2$ $y = -x - 7$ OR/OF $y = -x + c$ Subst./ <i>Vervang</i> $(-5; -2)$ $-2 = -(-5) + c$ $-7 = c$ $\therefore y = -x - 7$	✓ method / <i>metode</i> ✓ answer / <i>antwoord</i> Note: Neem kennis ✓✓ Answer only – Full marks <i>Slegs antwoord - Volpunte</i> (2)
		[11]


QUESTION 5/VRAAG 5

		
5.1	$x = \frac{-b}{2a}$ $x = \frac{-(-5)}{2(1)}$ $x = \frac{5}{2} = 2,5$ OR / OF $f(x) = x^2 - 5x + 6$ $f'(x) = 2x - 5$ $2x - 5 = 0$ $2x = 5$ $\therefore x = \frac{5}{2} = 2,5$	✓ substitution / <i>vervang</i> ✓ equation / <i>vergelyking</i> OR / OF ✓ $f'(x) = 0$ ✓ equation / <i>vergelyking</i> (2)
5.2	$f(x) = g(x)$ $x^2 - 5x + 6 = x + 1$ $x^2 - 6x + 5 = 0$ $(x - 5)(x - 1) = 0$ $x = 5 \text{ or/ of}$ $g(5) = 5 + 1 = 6$ $g(1) = 1 + 1 = 2$ B(1;2) and/en C(5;6)	✓ $f(x) = g(x)$ ✓ standard form / <i>standaardvorm</i> ✓ x-values / <i>x-waardes</i> ✓ y-values / <i>y-waardes</i> (4)




QUESTION 6/VRAAG 6

		
6.1	$f(x) = -\log_c x$ $\frac{1}{2} = -\log_c \left(\frac{1}{2}\right)$ $\sqrt{c} = 2$ $c = 4$ $g(x) = d x^2$ $\frac{1}{2} = d \left(\frac{1}{2}\right)^2$ $d = 2$	✓ subst. of / vervanging van $\left(\frac{1}{2}; \frac{1}{2}\right)$ ✓ value of c / waarde van c ✓ value of d / waarde van d (3)
6.2.1	$g: y = 2x^2$ $g^{-1}: x = 2y^2$ $g^{-1}: y^2 = \frac{1}{2}x$ $g^{-1}: y = \pm \sqrt{\frac{1}{2}x}$ $\therefore y = \sqrt{\frac{1}{2}x}$	✓ swopping x and y omruil van x en y ✓ answer / antwoord (2)
6.2.2	$f(x) = -\log_4 x$ $h(x) = \log_4 x$ $h^{-1}(x): y = 4^x$	✓ $h(x) = \log_4 x$ ✓ answer / antwoord (2)
6.2.3	$x \in \square$	✓ answer / antwoord (1)
		[8]

QUESTION 7/VRAAG 7

7.1	$A = P(1-i)^n$ $= 180000(1-0,13)^6$ $= R78\,052,72$	✓ $n = 6$ ✓ substitution / vervanging ✓ answer / antwoord (3)
7.2	$F_v = \frac{x[(1+i)^n - 1]}{i}$ $= \frac{900 \left[\left(1 + \frac{0,08}{12}\right)^{120} - 1 \right]}{\frac{0,08}{12}} + \frac{1300 \left[\left(1 + \frac{0,08}{12}\right)^{60} - 1 \right]}{\frac{0,08}{12}}$ $= 164\,651,4317 + 95\,519,91312$ $= R260\,171,34$	✓ $n = 120$ and/en $i = \frac{8\%}{12}$ or / of $\frac{8}{1200}$ ✓ $n = 60$ in F ✓ substitution into F_v / vervanging in F_v ✓ answer / antwoord (5)
7.3.1	$OB = P(1+i)^n - \frac{x[(1+i)^n - 1]}{i}$ $OB = 850000 \left(1 + \frac{0,13}{12}\right)^{75} - \frac{9958,39 \left[\left(1 + \frac{0,13}{12}\right)^{75} - 1 \right]}{\frac{0,13}{12}}$ $= R\,763\,890,54$ OR/OF $OB = \frac{x[1 - (1+i)^{-n}]}{i}$ $OB = \frac{9958,39 \left[1 - \left(1 + \frac{0,13}{12}\right)^{-165} \right]}{\frac{0,13}{12}}$ $= R\,763\,889,86$	✓ correct substitution into A formula / korrekte vervanging in A formule ✓ correct substitution into Fv formula / korrekte vervanging in Fv formule ✓ answer / antwoord OR/OF ✓ $n = 165$ ✓ substitution into the correct formula / vervanging in korrekte formule ✓ answer / antwoord (3)

7.3.2

**For Outstanding Balance / Vir Uitstaande Balans** $= R\ 763\ 890,54 :$

$$A = P(1+i)^n$$

$$A = 763890,54 \left(1 + \frac{0,13}{12}\right)^4$$

$$A = R\ 797\ 534,2651$$

$$P = \frac{x \left[1 - \left(1 + \frac{i}{12}\right)^{-n}\right]}{i}$$

$$797534,2651 = \frac{x \left[1 - \left(1 + \frac{0,13}{12}\right)^{-161}\right]}{\frac{0,13}{12}}$$

$$x = \frac{797534,2651 \times \frac{0,13}{12}}{\left[1 - \left(1 + \frac{0,13}{12}\right)^{-161}\right]}$$

$$x = R\ 10\ 490,96$$

Adjusted instalment is R10 490,96

OR / OF**For Outstanding Balance / Vir Uitstaande Balans** $= R\ 763\ 889,86 :$

$$A = P(1+i)^n$$

$$A = 763889,86 \left(1 + \frac{0,13}{12}\right)^4$$

$$= R\ 797\ 533,5551$$

$$797533,56 = \frac{x \left[1 - \left(1 + \frac{0,13}{12}\right)^{-161}\right]}{\frac{0,13}{12}}$$

$$x = \frac{797533,56 \times \frac{0,13}{12}}{\left[1 - \left(1 + \frac{0,13}{12}\right)^{-161}\right]}$$

$$x = R\ 10\ 490,95$$

✓ substitution into the correct A formula / vervanging in korrekte A formule

✓ accumulated amount / opgeboude bedrag

✓ substitution into the correct formula / vervanging in korrekte formule

✓ $n = -161$

✓ answer / antwoord

(5)

OR / OF

✓ substitution into the correct A formula / vervanging in korrekte A formule

✓ accumulated amount/opgeboude bedrag

✓ substitution into the correct formula / vervanging in korrekte formule

✓ $n = -161$

✓ answer / antwoord

(5)

[16]


QUESTION 8/VRAAG 8

8.1	$f(x) = x^2 - 3$ $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $= \lim_{h \rightarrow 0} \frac{(x+h)^2 - 3 - (x^2 - 3)}{h}$ $= \lim_{h \rightarrow 0} \frac{x^2 + 2xh - h^2 - 3 - (x^2 - 3)}{h}$ $= \lim_{h \rightarrow 0} \frac{x^2 + 2xh - h^2 - 3 - x^2 + 3}{h}$ $= \lim_{h \rightarrow 0} \frac{2xh - h^2}{h}$ $= \lim_{h \rightarrow 0} \frac{h(2x - h)}{h}$ $= \lim_{h \rightarrow 0} 2x - h$ $f'(x) = 2x$	✓ substitution into the formula <i>vervang in die formule</i> ✓ simplification / <i>vereenvoudiging</i> ✓ factorisation / <i>faktorisering</i> ✓ answer / <i>antwoord</i> (4)
8.2.1	$\frac{dy}{dx} = -6x + 7$	✓ $-6x$ ✓ 7 (2)
8.2.2	$D_x \left[\frac{x^3 - 5x^2}{x^3} - \sqrt{x} \right]$ $= D_x \left[\frac{x^3}{x^3} - \frac{5x^2}{x^3} - x^{\frac{1}{2}} \right]$ $= D_x \left[1 - 5x^{-1} - x^{\frac{1}{2}} \right]$ $= 0 + 5x^{-2} - \frac{1}{2}x^{-\frac{1}{2}}$ $= \frac{5}{x^2} - \frac{1}{2\sqrt{x}}$	✓ $x^{\frac{1}{2}}$ ✓ $1 - 5x^{-1}$ ✓ 0 & $5x^{-2}$ (zero does not have to be seen) (<i>hoef nie nul te sien nie</i>) ✓ $-\frac{1}{2}x^{-\frac{1}{2}}$ (4)

(3)
[13]





QUESTION 9/VRAAG 9



9.1.1	$f(x) = -3x^3 + mx^2 + nx$ $f'(x) = -9x^2 + 2mx + n$ $0 = -9(1)^2 + 2m(1) + n$ $0 = -9 + 2m + n$ $n = 9 - 2m \dots \dots \dots (1)$ $f'(x) = -9x^2 + 2mx + n$ $0 = -9(3)^2 + 2m(3) + n$ $81 = 6m + n$ $81 - 6m = n \dots \dots \dots (2)$ $81 - 6m = 9 - 2m$ $81 - 9 = 6m - 2m$ $72 = 4m$ $\therefore m = 18$ $n = 9 - 2(18)$ $\therefore n = -27$ $\therefore f(x) = -3x^3 + 18x^2 - 27x$	✓ equation 1 / vergelyking 1 ✓ equation 2 / vergelyking 2 ✓ equating (method) / gelykstel (metode) ✓ solve for m / oplos vir m ✓ substituting value of m / vervanging van waarde van m (5)
9.1.2	$f(a)$ – corresponding y-value when $x = a$, while $f'(a)$ – gradient/derivative/rate of change of when $x = a$ $f(a)$ – is die ooreenstemmende y-waarde wanneer $x = a$, terwyl $f'(a)$ – stel voor die gradiënt/afgeleide/ veranderingskoers van f wanneer $x = a$	explanation of / verduideliking van ✓ $f(a)$ ✓ $f'(a)$ (2)

9.1.3	Point of inflection/<i>Buigpunt/infleksiepunt</i> $x = \frac{x_A + x_B}{2}$ $x = \frac{1+3}{2} = 2$ OR / OF $f'(x) = -9x^2 + 36x - 27$ $f''(x) = -18x + 36$ $f''(x) = 0$ $-18x + 36 = 0$ $18x = 36$ $\therefore x = 2$ $\therefore f(2) = -3(2)^3 + 18(2)^2 - 27(2) = -6$ $\therefore (2; -6)$ Gradient of / <i>Gradiënt van</i> $g(x)$: $f'(2) = -9(2)^2 + 36(2) - 27 = 9$ Gradient of / <i>Gradiënt van</i> $h(x)$: $m_g \times m_h = -1$ $\therefore m_h = -\frac{1}{9}$ $\therefore h(x) = -\frac{1}{9}x$	✓ method / <i>metode</i> ✓ ✓ $f(2)$ ✓ $f'(2)$ ✓ $h(x) = -\frac{1}{9}x$ (5)
9.1.4	$f''(x) > 0$ when $f(x)$ is concave up x – value for point of inflection is 2 $\therefore x < 2$ $f''(x) > 0$ wanneer $f(x)$ konkaaf op is x-waarde vir buigpunt/infleksiepunt is 2 $\therefore x < 2$	✓✓ answer / <i>antwoord</i> (2)

9.2	• $a = 2 > 0$  • $(-3;0)$ $(0;0)$ $(3;0)$ (x-intercepts) (x-afsnitte)  • x-values of stationary points: x-waardes van stasionêre punte 	✓ shape / vorm ✓ intercepts on the graph/ afsnitte op die grafiek ✓ x- values for stationary points x-waardes vir stasionêre punte (3) [17]
------------	--	--





QUESTION 10/VRAAG 10

10.1	$S(t) = -3t^2 + 30t$ $S(3) = -3(3)^2 + 30(3)$ $= 63$ scripts / skrifte	✓ substitution / <i>vervanging</i> ✓ answer / <i>antwoord</i> (2)																				
10.2	$S'(t) = -6t + 30$ For maximum number of scripts, $S'(t) = 0$ / <i>Vir maksimum aantal skrifte, $S'(t) = 0$</i> $-6t + 30 = 0$ $6t = 30$ $t = 5$ (Day 5 / <i>Dag 5</i>)	✓ $S'(t)$ ✓ $S'(t) = 0$ ✓ $t = 5$ (3)																				
10.3	No / <i>Nee</i> <table border="1"><tr><td>D1</td><td>D2</td><td>D3</td><td>D4</td><td>D5</td><td>D6</td><td>D7</td><td>D8</td><td>D9</td><td>D10</td></tr><tr><td>27</td><td>48</td><td>63</td><td>72</td><td>75</td><td>72</td><td>63</td><td>48</td><td>27</td><td>0</td></tr></table> Sum/ <i>Som</i> = 495 scripts/ <i>skrifte</i>	D1	D2	D3	D4	D5	D6	D7	D8	D9	D10	27	48	63	72	75	72	63	48	27	0	✓ No / <i>Nee</i> ✓ explanation / <i>verduideliking</i> (2)
D1	D2	D3	D4	D5	D6	D7	D8	D9	D10													
27	48	63	72	75	72	63	48	27	0													
		[7]																				



QUESTION 11/VRAAG 11

		
11.1.1	$x = 1 - (0,11 + 0,19 + 0,41)$ $x = 0,29$ $P(A) = 0,29 + 0,11 = 0,4$	✓ value of x / waarde van x ✓ answer / antwoord (2)
11.1.2	$P(A \text{ or/of not/nie } B) = 0,29 + 0,11 + 0,41 = 0,81$	✓✓ answer / antwoord (2)
11.2.1	$a = 4$	✓ answer / antwoord (1)
11.2.2	$\frac{14}{30} = \frac{7}{15}$	✓ answer / antwoord (1)
11.2.3	$P(\text{winning a game}) = \frac{7}{30}$ $P(\text{playing at home}) = \frac{15}{30} = \frac{1}{2}$ $P(\text{winning a game}) \times P(\text{playing at home})$ $= \frac{7}{30} \times \frac{1}{2}$ $= \frac{7}{60} = 0,12$ $P(\text{winning a game and playing at home}) = \frac{3}{30} = 0,10$ events are not independent, since $P(\text{winning a game and playing at home}) \neq$ $P(\text{winning a game}) \times P(\text{playing at home})$	✓ $P(\text{winning a game}) \times P(\text{playing at home})$ ✓ $P(\text{winning a game and playing at home})$ ✓ conclusion
	$P(\text{wen 'n wedstryd}) = \frac{7}{30}$ $P(\text{speel tuiswedstryd}) = \frac{15}{30} = \frac{1}{2}$ $P(\text{wen wedstryd}) \times P(\text{speel tuiswedstryd})$ $= \frac{7}{30} \times \frac{1}{2}$ $= \frac{7}{60} = 0,12$ $P(\text{wen wedstryd en tuis wedstryd}) = \frac{3}{30} = 0,10$ gebeurtenisse is nie onafhanklik nie, omdat $P(\text{wen wedstryd en speel tuiswedstryd}) \neq$ $P(\text{wen wedstryd}) \times P(\text{speel tuiswedstryd})$	 ✓ $P(\text{wen wedstryd}) \times P(\text{speel tuiswedstryd})$ ✓ $P(\text{wen wedstryd en speel tuiswedstryd})$ ✓ Gevolgtrekking (3)
		[9]

QUESTION 12/VRAAG 12

12.1	$21 \times 20 \times 10 \times 9 \times 19 \times 18$ $= 12\,927\,600$ codes/kodes	✓✓ answer / antwoord (2)																									
12.2	DIGITS / SYFERS: <table> <tr> <td><u>4</u></td> <td><u>2</u></td> <td>or/of</td> <td><u>3</u></td> <td><u>1</u></td> </tr> <tr> <td>4</td> <td>1</td> <td></td> <td>4</td> <td>1</td> </tr> <tr> <td>6</td> <td>2</td> <td></td> <td>6</td> <td>2</td> </tr> <tr> <td>8</td> <td>4</td> <td></td> <td>8</td> <td>4</td> </tr> <tr> <td>9</td> <td></td> <td></td> <td>9</td> <td></td> </tr> </table>  LETTERS BEFORE G / LETTERS VOOR G: A; B; C; D; E and/en F Out of 6 letters remove A and E (vowels) = 4 letters <i>Van die 6 letters verwyder A en E (klinkers) = 4 letters</i> COMBINED / KOMBINASIE : $n(A) = (4 \times 20 \times 4 \times 2 \times 19 \times 18) + (4 \times 20 \times 3 \times 1 \times 19 \times 18)$ $= 218\,880 + 82\,080$ $= 300\,960$ $P(A) = \frac{n(A)}{n(S)}$ $= \frac{300960}{12927600}$ $= \frac{22}{945}$ $\approx 0,02$	<u>4</u>	<u>2</u>	or/of	<u>3</u>	<u>1</u>	4	1		4	1	6	2		6	2	8	4		8	4	9			9		✓ 4 in $n(A)$ ✓ 4×2 and/en 3×1 in $n(A)$ ✓ dividing by / deel deur 12927600 ✓ answer / antwoord (4) [6]
<u>4</u>	<u>2</u>	or/of	<u>3</u>	<u>1</u>																							
4	1		4	1																							
6	2		6	2																							
8	4		8	4																							
9			9																								

TOTAL/TOTAAL: 150